

# POSITIVE SOLUTIONS OF A NONLINEAR THREE-POINT EIGENVALUE PROBLEM WITH INTEGRAL BOUNDARY CONDITIONS

FAOUZI HADDOUCHI AND SLIMANE BENAICHA

ABSTRACT. In this paper, we study the existence of positive solutions of a three-point integral boundary value problem (BVP) for the following second-order differential equation

$$\begin{aligned}u''(t) + \lambda a(t)f(u(t)) &= 0, \quad 0 < t < 1, \\u'(0) = 0, \quad u(1) &= \alpha \int_0^\eta u(s)ds,\end{aligned}$$

where  $\lambda > 0$  is a parameter,  $0 < \eta < 1$ ,  $0 < \alpha < \frac{1}{\eta}$ . By using the properties of the Green's function and Krasnoselskii's fixed point theorem on cones, the eigenvalue intervals of the nonlinear boundary value problem are considered, some sufficient conditions for the existence of at least one positive solutions are established.

**Mathematics Subject Classification (2010):** 34B15, 34C25, 34B18

**Key words:** Positive solutions, Krasnoselskii's fixed point theorem, Three-point integral boundary value problems, Eigenvalue, Cone.

*Article history:*

Received 16 September 2015

Received in revised form 26 December 2015

Accepted 27 December 2015

## 1. INTRODUCTION

In this work, we study the existence of positive solutions of a three-point integral boundary value problem (BVP) for the following second-order differential equation:

$$(1.1) \quad u''(t) + \lambda a(t)f(u(t)) = 0, \quad t \in (0, 1),$$

$$(1.2) \quad u'(0) = 0, \quad u(1) = \alpha \int_0^\eta u(s)ds,$$

where  $0 < \eta < 1$  and  $0 < \alpha < \frac{1}{\eta}$ ,  $\lambda$  is a positive parameter, and

(H1)  $f \in C([0, \infty), [0, \infty))$ ;

(H2)  $a \in C([0, 1], [0, \infty))$  and there exists  $t_0 \in [0, \eta]$  such that  $a(t_0) > 0$ .

The study of the existence of solutions of multi-point boundary value problems for linear second-order ordinary differential equations was initiated by Il'in and Moiseev [12]. Then Gupta [8] studied three-point boundary value problems for nonlinear second-order ordinary differential equations. Since then, the existence of positive solutions for nonlinear second order three-point boundary-value problems has been studied by many authors by using the fixed point theorem, nonlinear alternative of the Leray-Schauder

approach, or coincidence degree theory. We refer the reader to [1], [2], [4], [6], [7], [10], [11], [17]-[27], [29], [30], [32]-[36] and the references therein.

Liu [20] proved the existence of single and multiple positive solutions for the three-point boundary value problem (BVP)

$$(1.3) \quad u''(t) + a(t)f(u(t)) = 0, \quad t \in (0, 1),$$

$$(1.4) \quad u'(0) = 0, \quad u(1) = \beta u(\eta),$$

where  $0 < \eta < 1$  and  $0 < \beta < 1$ .

Recently, Ma [28] studied the second-order three-point boundary value problem (BVP)

$$(1.5) \quad u''(t) + \lambda a(t)f(u(t)) = 0, \quad t \in (0, 1),$$

$$(1.6) \quad u(0) = \beta u(\eta), \quad u(1) = \alpha u(\eta),$$

where  $\lambda > 0$  is a parameter,  $\alpha > 0$ ,  $\beta > 0$ ,  $0 < \eta < 1$ ,  $a \in C([0, 1], [0, \infty))$ ,  $f \in C([0, \infty), [0, \infty))$  and there exists  $x_0 \in (0, 1)$  such that  $a(x_0) > 0$ . She obtained the existence of single and multiple positive solutions by using Krasnoselskii's fixed point theorem in cones [16].

Boundary value problems with integral boundary conditions for ordinary differential equations represent a very interesting and important class of problems, and arise in the study of various physical, biological and chemical processes, such as heat conduction, chemical engineering, underground water flow, thermo-elasticity, and plasma physics. They include two, three, multi-point and nonlocal BVPs as special cases. The existence of positive solutions for such class of problems has attracted much attention (see [3], [5], [13]-[15], [31], [34], [37] and the references therein).

In [34], Tariboon and Sitthiwiratham investigated the existence of positive solutions of the following three-point integral boundary value problem (BVP)

$$(1.7) \quad u''(t) + a(t)f(u(t)) = 0, \quad t \in (0, 1),$$

$$(1.8) \quad u(0) = 0, \quad u(1) = \alpha \int_0^\eta u(s)ds,$$

where  $0 < \eta < 1$  and  $0 < \alpha < \frac{2}{\eta^2}$ ,  $f \in C([0, \infty), [0, \infty))$ ,  $a \in C([0, 1], [0, \infty))$  and there exists  $t_0 \in [\eta, 1]$  such that  $a(t_0) > 0$ . They showed the existence of at least one positive solution if  $f$  is either superlinear or sublinear by applying Krasnoselskii's fixed point theorem in cones [16].

In [9], by using Leggett-Williams fixed-point theorem, the authors considered the multiplicity of positive solutions of the following three-point integral boundary value problem (BVP)

$$(1.9) \quad u''(t) + f(t, u(t)) = 0, \quad t \in (0, T),$$

$$(1.10) \quad u(0) = \beta u(\eta), \quad u(T) = \alpha \int_0^\eta u(s)ds,$$

where  $0 < \eta < T$ ,  $0 < \alpha < \frac{2T}{\eta^2}$ ,  $0 \leq \beta < \frac{2T - \alpha\eta^2}{\alpha\eta^2 - 2\eta + 2T}$ , and  $f \in C([0, T] \times [0, \infty), [0, \infty))$ .

Motivated greatly by the above-mentioned excellent works, the aim of this paper is to establish some sufficient conditions for the existence of at least one positive solutions of the BVP (1.1) and (1.2). Our ideas are similar those used in [28], but a little different.

We firstly give the corresponding Green's function for the associated linear BVP and some of its properties. Moreover, by applying Krasnoselskii's fixed point theorem, we derive an interval of  $\lambda$  on which there exists a positive solution for the three-point integral boundary value problem (1.1) and (1.2).

As applications, some interesting examples are presented to illustrate the main results. The key tool in our approach is the following Krasnoselskii's fixed point theorem in a cone [16].

**Theorem 1.1.** [16]. *Let  $E$  be a Banach space, and let  $K \subset E$  be a cone. Assume that  $\Omega_1, \Omega_2$  are open bounded subsets of  $E$  with  $0 \in \Omega_1, \bar{\Omega}_1 \subset \Omega_2$ , and let*

$$A : K \cap (\bar{\Omega}_2 \setminus \Omega_1) \longrightarrow K$$

*be a completely continuous operator such that either*

*(i)  $\|Au\| \leq \|u\|$ ,  $u \in K \cap \partial\Omega_1$ , and  $\|Au\| \geq \|u\|$ ,  $u \in K \cap \partial\Omega_2$ ; or*

*(ii)  $\|Au\| \geq \|u\|$ ,  $u \in K \cap \partial\Omega_1$ , and  $\|Au\| \leq \|u\|$ ,  $u \in K \cap \partial\Omega_2$*

*hold. Then  $A$  has a fixed point in  $K \cap (\bar{\Omega}_2 \setminus \Omega_1)$ .*

## 2. PRELIMINARIES

**Lemma 2.1.** *Let  $\alpha\eta \neq 1$ . Then for  $y \in C([0, 1], \mathbb{R})$ , the problem*

$$(2.1) \quad u''(t) + y(t) = 0, \quad t \in (0, 1),$$

$$(2.2) \quad u'(0) = 0, \quad u(1) = \alpha \int_0^\eta u(s) ds,$$

*has a unique solution*

$$(2.3) \quad u(t) = \int_0^1 G(t, s) y(s) ds,$$

*where  $G(t, s) : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$  is the Green's function defined by*

$$(2.4) \quad G(t, s) = \frac{1}{2(1 - \alpha\eta)} \begin{cases} 2(1 - s) - \alpha(\eta - s)^2 - 2(1 - \alpha\eta)(t - s), & s \leq \min\{\eta, t\}; \\ 2(1 - s) - \alpha(\eta - s)^2, & t \leq s \leq \eta; \\ 2(1 - s) - 2(1 - \alpha\eta)(t - s), & \eta \leq s \leq t; \\ 2(1 - s), & \max\{\eta, t\} \leq s. \end{cases}$$

*Proof.* From (2.1), we have

$$(2.5) \quad u(t) = u(0) - \int_0^t (t - s) y(s) ds.$$

Integrating (2.5) from 0 to  $\eta$ , where  $\eta \in (0, 1)$ , we have

$$\int_0^\eta u(s) ds = u(0)\eta - \frac{1}{2} \int_0^\eta (\eta - s)^2 y(s) ds.$$

Since

$$u(1) = u(0) - \int_0^1 (1 - s) y(s) ds,$$

from  $u(1) = \alpha \int_0^\eta u(s) ds$ , we have

$$(1 - \alpha\eta)u(0) = \int_0^1 (1 - s) y(s) ds - \frac{\alpha}{2} \int_0^\eta (\eta - s)^2 y(s) ds.$$

Therefore,

$$u(0) = \frac{1}{1 - \alpha\eta} \int_0^1 (1 - s) y(s) ds - \frac{\alpha}{2(1 - \alpha\eta)} \int_0^\eta (\eta - s)^2 y(s) ds,$$

from which it follows that

$$\begin{aligned} u(t) &= \frac{1}{1 - \alpha\eta} \int_0^1 (1 - s) y(s) ds - \frac{\alpha}{2(1 - \alpha\eta)} \int_0^\eta (\eta - s)^2 y(s) ds \\ &\quad - \int_0^t (t - s) y(s) ds. \end{aligned}$$

If  $t \leq \eta$ , then

$$\begin{aligned}
u(t) &= \int_0^\eta \frac{1-s}{1-\alpha\eta} y(s) ds - \int_0^\eta \frac{\alpha(\eta-s)^2}{2(1-\alpha\eta)} y(s) ds + \int_\eta^1 \frac{1-s}{1-\alpha\eta} y(s) ds \\
&\quad - \int_0^t (t-s)y(s) ds \\
&= \int_0^t \frac{2(1-s) - \alpha(\eta-s)^2 - 2(1-\alpha\eta)(t-s)}{2(1-\alpha\eta)} y(s) ds + \int_\eta^1 \frac{1-s}{1-\alpha\eta} y(s) ds \\
&\quad + \int_t^\eta \frac{2(1-s) - \alpha(\eta-s)^2}{2(1-\alpha\eta)} y(s) ds \\
&= \int_0^1 G(t,s)y(s) ds.
\end{aligned}$$

If  $t \geq \eta$ , then

$$\begin{aligned}
u(t) &= \int_0^t \frac{1-s}{1-\alpha\eta} y(s) ds - \int_0^\eta \frac{\alpha(\eta-s)^2}{2(1-\alpha\eta)} y(s) ds + \int_t^1 \frac{1-s}{1-\alpha\eta} y(s) ds \\
&\quad - \int_0^t (t-s)y(s) ds \\
&= \int_0^\eta \frac{2(1-s) - \alpha(\eta-s)^2 - 2(1-\alpha\eta)(t-s)}{2(1-\alpha\eta)} y(s) ds + \int_t^1 \frac{1-s}{1-\alpha\eta} y(s) ds \\
&\quad + \int_\eta^t \frac{(1-s) - (1-\alpha\eta)(t-s)}{1-\alpha\eta} y(s) ds \\
&= \int_0^1 G(t,s)y(s) ds.
\end{aligned}$$

This completes the proof. □

For convenience, we define

$$g(s) = \frac{1}{1-\alpha\eta}(1-s), \quad s \in [0, 1].$$

For the Green's function  $G(t, s)$ , we have the following two lemmas.

**Lemma 2.2.** *Let  $0 < \eta < 1$  and  $0 < \alpha < \frac{1}{\eta}$ . Then the Green's function in (2.4) satisfies*

$$(2.6) \quad 0 \leq G(t, s) \leq g(s),$$

for each  $s, t \in [0, 1]$ .

*Proof.* First of all, note that by (2.4) it follows that  $G(t, s) \leq g(s)$  for any  $(t, s) \in [0, 1] \times [0, 1]$ .

Next, we will prove that  $G(t, s) \geq 0$  for any  $(t, s) \in [0, 1] \times [0, 1]$ .

If  $\eta \leq s \leq t$ , then

$$\begin{aligned}
G(t, s) &= \frac{1}{1-\alpha\eta} \left( (1-s) - (1-\alpha\eta)(t-s) \right) \\
&= \frac{1}{1-\alpha\eta} \left( (1-t) + \alpha\eta(t-s) \right) \geq 0.
\end{aligned}$$

If  $t \leq s \leq \eta$ , then

$$\begin{aligned}
G(t, s) &= \frac{1}{2(1-\alpha\eta)} \left( 2(1-s) - \alpha(\eta-s)^2 \right) \\
&\geq \frac{1}{2(1-\alpha\eta)} \left( 2(1-s) - \frac{(\eta-s)^2}{\eta} \right) \\
&= \frac{2\eta - \eta^2 - s^2}{2\eta(1-\alpha\eta)} \\
&\geq \frac{2\eta - 2\eta^2}{2\eta(1-\alpha\eta)} \\
&= \frac{1-\eta}{1-\alpha\eta} \\
&\geq 0.
\end{aligned}$$

If  $s \leq \min\{\eta, t\}$ , then

$$\begin{aligned}
G(t, s) &= \frac{1}{2(1-\alpha\eta)} \left( 2(1-s) - \alpha(\eta-s)^2 - 2(1-\alpha\eta)(t-s) \right) \\
&= \frac{1}{2(1-\alpha\eta)} \left( 2(1-t) - \alpha\eta^2 - \alpha s^2 + 2\alpha\eta t \right) \\
&\geq \frac{1}{2(1-\alpha\eta)} \left( 2(1-t) + \alpha\eta(t-\eta) + \alpha s(\eta-s) \right).
\end{aligned}$$

We distinguish the following two cases:

If  $s \leq \eta \leq t$ , then  $G(t, s) \geq 0$ .

If  $s \leq t \leq \eta$ , then

$$\begin{aligned}
G(t, s) &= \frac{1}{2(1-\alpha\eta)} \left( 2(1-t) - \alpha\eta^2 - \alpha s^2 + 2\alpha\eta t \right) \\
&\geq \frac{1}{2(1-\alpha\eta)} \left( 2(1-t) - \alpha(\eta-t)^2 \right) \\
&\geq \frac{1}{2(1-\alpha\eta)} \left( 2(1-t) - \frac{(\eta-t)^2}{\eta} \right) \\
&= \frac{2\eta - \eta^2 - t^2}{2\eta(1-\alpha\eta)} \\
&\geq \frac{2\eta - 2\eta^2}{2\eta(1-\alpha\eta)} \\
&= \frac{1-\eta}{1-\alpha\eta} \\
&\geq 0.
\end{aligned}$$

The proof is completed. □

**Lemma 2.3.** Let  $0 < \eta < 1$  and  $0 < \alpha < \frac{1}{\eta}$ . Then for any  $(t, s) \in [0, \eta] \times [0, 1]$ ,  $G(t, s) \geq \gamma g(s)$ , where

$$(2.7) \quad 0 < \gamma = 1 - \eta < 1.$$

*Proof.* If  $s = 1$ , then by Lemma 2.2, the result follows. Now we suppose that  $(t, s) \in [0, \eta] \times [0, 1)$ .

If  $t \leq s \leq \eta$ , then

$$\begin{aligned}
\frac{G(t,s)}{g(s)} &= \frac{2(1-s) - \alpha(\eta-s)^2}{2(1-s)} \\
&\geq \frac{2(1-s) - \frac{(\eta-s)^2}{\eta}}{2(1-s)} \\
&= \frac{2\eta - \eta^2 - s^2}{2\eta(1-s)} \\
&\geq \frac{1-\eta}{1-s} \\
&\geq \gamma.
\end{aligned}$$

If  $s \leq t \leq \eta$ , then

$$\begin{aligned}
\frac{G(t,s)}{g(s)} &= \frac{2(1-s) - \alpha(\eta-s)^2 - 2(1-\alpha\eta)(t-s)}{2(1-s)} \\
&= \frac{2(1-t) - \alpha\eta^2 - \alpha s^2 + 2\alpha\eta t}{2(1-s)} \\
&\geq \frac{2(1-t) - \alpha(\eta-t)^2}{2\eta(1-s)} \\
&\geq \frac{1-\eta}{1-s} \\
&\geq \gamma.
\end{aligned}$$

If  $t \leq \eta \leq s$ , then

$$\frac{G(t,s)}{g(s)} = 1 \geq \gamma.$$

Therefore,

$$G(t,s) \geq \gamma g(s), \quad (t,s) \in [0,\eta] \times [0,1].$$

The proof is completed. □

Let  $E = C([0,1], \mathbb{R})$ , and only the sup norm is used. It is easy to see that the BVP (1.1) and (1.2) has a solution  $u = u(t)$  if and only if  $u$  is a fixed point of operator  $A_\lambda$ , where  $A_\lambda$  is defined by

$$\begin{aligned}
A_\lambda u(t) &= \frac{\lambda}{1-\alpha\eta} \int_0^1 (1-s)a(s)f(u(s))ds - \frac{\lambda\alpha}{2(1-\alpha\eta)} \int_0^\eta (\eta-s)^2 a(s)f(u(s))ds \\
&\quad - \lambda \int_0^t (t-s)a(s)f(u(s))ds \\
&= \lambda \int_0^1 G(t,s)a(s)f(u(s))ds.
\end{aligned}$$

Denote

$$(2.8) \quad K = \left\{ u \in E : u \geq 0, \min_{t \in [0,\eta]} u(t) \geq \gamma \|u\| \right\},$$

where  $\gamma$  is defined in (2.7). It is obvious that  $K$  is a cone in  $E$ .

**Lemma 2.4.** *Assume that (H1) and (H2) hold,  $0 < \eta < 1$  and  $0 < \alpha < \frac{1}{\eta}$ . Then the operator  $A_\lambda : K \rightarrow K$  is completely continuous.*

*Proof.* For  $u \in K$ , according to the definition of  $A_\lambda$ , Lemma 2.2 and Lemma 2.3, it is easy to prove that  $A_\lambda K \subset K$ . By the Ascoli-Arzelà theorem, it is easy to show that  $A_\lambda : K \rightarrow K$  is completely continuous.  $\square$

In what follows, for the sake of convenience, set

$$\Lambda_1 = \frac{1}{1 - \alpha\eta} \int_0^1 (1 - s)a(s)ds, \quad \Lambda_2 = \frac{\gamma}{2(1 - \alpha\eta)} \int_0^\eta \left(2(1 - \eta) + \alpha(\eta^2 - s^2)\right)a(s)ds,$$

$$f_0 = \lim_{u \rightarrow 0^+} \frac{f(u)}{u}, \quad f_\infty = \lim_{u \rightarrow \infty} \frac{f(u)}{u}.$$

### 3. MAIN RESULTS

In this section, we will state and prove our main results.

**Theorem 3.1.** *Suppose that (H1) and (H2) hold,  $0 < \eta < 1$  and  $0 < \alpha < \frac{1}{\eta}$ . If  $\Lambda_1 f_0 < \Lambda_2 f_\infty$ , then for each  $\lambda \in (\frac{1}{\Lambda_2 f_\infty}, \frac{1}{\Lambda_1 f_0})$ , the BVP (1.1) and (1.2) has at least one positive solution.*

*Proof.* Let  $\lambda \in (\frac{1}{\Lambda_2 f_\infty}, \frac{1}{\Lambda_1 f_0})$ , and choose  $\epsilon > 0$  such that

$$(3.1) \quad \frac{1}{\Lambda_2(f_\infty - \epsilon)} \leq \lambda \leq \frac{1}{\Lambda_1(f_0 + \epsilon)}.$$

By the definition of  $f_0$ , there exists  $\rho_1 > 0$  such that

$$(3.2) \quad f(u) \leq (f_0 + \epsilon)u, \text{ for } u \in (0, \rho_1].$$

Let  $\Omega_{\rho_1} = \{u \in E : \|u\| < \rho_1\}$ , then from (3.1), (3.2) and Lemma 2.2, for any  $u \in K \cap \partial\Omega_{\rho_1}$ , we have

$$\begin{aligned} A_\lambda u(t) &= \lambda \int_0^1 G(t, s)a(s)f(u(s))ds \\ &\leq \lambda \int_0^1 g(s)a(s)f(u(s))ds \\ &= \frac{\lambda}{1 - \alpha\eta} \int_0^1 (1 - s)a(s)f(u(s))ds \\ &\leq \frac{\lambda}{1 - \alpha\eta} \int_0^1 (1 - s)a(s)(f_0 + \epsilon)u(s)ds \\ &\leq \lambda\Lambda_1(f_0 + \epsilon)\|u\| \leq \|u\|, \end{aligned}$$

which yields

$$(3.3) \quad \|A_\lambda u\| \leq \|u\|, \text{ for } u \in K \cap \partial\Omega_{\rho_1}.$$

Further, by the definition of  $f_\infty$ , there exists  $\widehat{\rho}_2 > 0$  such that

$$(3.4) \quad f(u) \geq (f_\infty - \epsilon)u, \text{ for } u \in [\widehat{\rho}_2, \infty).$$

Now, set  $\rho_2 = \max\left\{2\rho_1, \frac{\widehat{\rho}_2}{\gamma}\right\}$  and  $\Omega_{\rho_2} = \{u \in E : \|u\| < \rho_2\}$ . Then  $u \in K \cap \partial\Omega_{\rho_2}$  implies that

$$u(t) \geq \gamma\|u\| \geq \widehat{\rho}_2, \quad t \in [0, \eta],$$

and so,

$$\begin{aligned}
A_\lambda u(\eta) &= \lambda \int_0^1 G(\eta, s) a(s) f(u(s)) ds \\
&\geq \lambda \int_0^\eta G(\eta, s) a(s) f(u(s)) ds \\
&\geq \lambda \int_0^\eta G(\eta, s) a(s) (f_\infty - \epsilon) u(s) ds \\
&\geq \lambda \gamma (f_\infty - \epsilon) \|u\| \int_0^\eta G(\eta, s) a(s) ds \\
&= \lambda \gamma (f_\infty - \epsilon) \|u\| \int_0^\eta \frac{2(1-s) - \alpha(\eta-s)^2 - 2(1-\alpha\eta)(\eta-s)}{2(1-\alpha\eta)} a(s) ds \\
&= \lambda (f_\infty - \epsilon) \|u\| \frac{\gamma}{2(1-\alpha\eta)} \int_0^\eta \left( 2(1-\eta) + \alpha(\eta^2 - s^2) \right) a(s) ds \\
&= \lambda \Lambda_2 (f_\infty - \epsilon) \|u\| \geq \|u\|.
\end{aligned}$$

This implies that

$$(3.5) \quad \|A_\lambda u\| \geq \|u\|, \quad \text{for } u \in K \cap \partial\Omega_{\rho_2}.$$

Therefore, from (3.3), (3.5) and Theorem 1.1, it follows that  $A_\lambda$  has a fixed point  $u$  with  $\rho_1 \leq \|u\| \leq \rho_2$  in  $K \cap (\overline{\Omega}_{\rho_2} \setminus \Omega_{\rho_1})$ , which is a desired positive solution of the BVP (1.1) and (1.2).  $\square$

By Theorem 3.1 we can easily obtain the following corollary.

**Corollary 3.2.** *Assume that (H1) and (H2) hold,  $0 < \eta < 1$  and  $0 < \alpha < \frac{1}{\eta}$ . Then we have*

- (1) *If  $f_0 = 0$ ,  $f_\infty = \infty$ , then for each  $\lambda \in (0, \infty)$ , the BVP (1.1) and (1.2) has at least one positive solution.*
- (2) *If  $f_\infty = \infty$ ,  $0 < f_0 < \infty$ , then for each  $\lambda \in (0, \frac{1}{\Lambda_1 f_0})$ , the BVP (1.1) and (1.2) has at least one positive solution.*
- (3) *If  $f_0 = 0$ ,  $0 < f_\infty < \infty$ , then for each  $\lambda \in (\frac{1}{\Lambda_2 f_\infty}, \infty)$ , the BVP (1.1) and (1.2) has at least one positive solution.*

**Theorem 3.3.** *Suppose that (H1) and (H2) hold,  $0 < \eta < 1$  and  $0 < \alpha < \frac{1}{\eta}$ . If  $\Lambda_1 f_\infty < \Lambda_2 f_0$ , then for each  $\lambda \in (\frac{1}{\Lambda_2 f_0}, \frac{1}{\Lambda_1 f_\infty})$ , the BVP (1.1) and (1.2) has at least one positive solution.*

*Proof.* Let  $\lambda \in (\frac{1}{\Lambda_2 f_0}, \frac{1}{\Lambda_1 f_\infty})$ , and choose  $\epsilon > 0$  such that

$$(3.6) \quad \frac{1}{\Lambda_2 (f_0 - \epsilon)} \leq \lambda \leq \frac{1}{\Lambda_1 (f_\infty + \epsilon)}.$$

By the definition of  $f_0$ , there exists  $\rho_1 > 0$  such that

$$(3.7) \quad f(u) \geq (f_0 - \epsilon)u, \quad \text{for } u \in (0, \rho_1].$$

Let  $\Omega_{\rho_1} = \{u \in E : \|u\| < \rho_1\}$ . Hence, for any  $u \in K \cap \partial\Omega_{\rho_1}$ , from (3.6), (3.7), we get

$$\begin{aligned}
A_\lambda u(\eta) &\geq \lambda \int_0^\eta G(\eta, s) a(s) f(u(s)) ds \\
&\geq \lambda \int_0^\eta G(\eta, s) a(s) (f_0 - \epsilon) u(s) ds \\
&\geq \lambda \Lambda_2 (f_0 - \epsilon) \|u\| \geq \|u\|.
\end{aligned}$$



Therefore

$$(3.8) \quad \|A_\lambda u\| \geq \|u\|, \quad \text{for } u \in K \cap \partial\Omega_{\rho_1}.$$

By the definition of  $f_\infty$ , there exists  $\rho_0 > 0$  such that

$$(3.9) \quad f(u) \leq (f_\infty + \epsilon)u, \quad \text{for } u \in [\rho_0, \infty).$$

Next, we consider two cases:

If  $f$  is bounded. Let  $f(u) \leq L$  for all  $u \in [0, \infty)$ . Set  $\rho_2 = \max\{2\rho_1, \lambda\Lambda_1 L\}$  and  $\Omega_{\rho_2} = \{u \in E : \|u\| < \rho_2\}$ , then for  $u \in K \cap \partial\Omega_{\rho_2}$ , we have

$$\begin{aligned} A_\lambda u(t) &\leq \frac{\lambda}{1 - \alpha\eta} \int_0^1 (1-s)a(s)f(u(s))ds \\ &\leq \lambda L \Lambda_1 \leq \rho_2 \leq \|u\|. \end{aligned}$$

Therefore

$$(3.10) \quad \|A_\lambda u\| \leq \|u\|, \quad \text{for } u \in K \cap \partial\Omega_{\rho_2}.$$

If  $f$  is unbounded, then from  $f \in C([0, \infty), [0, \infty))$ , we know that there is  $\rho_2: \rho_2 \geq \max\{2\rho_1, \gamma^{-1}\rho_0\}$  such that

$$(3.11) \quad f(u) \leq f(\rho_2), \quad \text{for } u \in [0, \rho_2].$$

Let  $\Omega_{\rho_2} = \{u \in E : \|u\| < \rho_2\}$ , then for  $u \in K \cap \partial\Omega_{\rho_2}$ , we have

$$\begin{aligned} A_\lambda u(t) &\leq \frac{\lambda}{1 - \alpha\eta} \int_0^1 (1-s)a(s)f(u(s))ds \\ &\leq \frac{\lambda}{1 - \alpha\eta} \int_0^1 (1-s)a(s)f(\rho_2)ds \\ &\leq \lambda\Lambda_1\rho_2(f_\infty + \epsilon) \leq \rho_2 = \|u\|. \end{aligned}$$

Thus

$$(3.12) \quad \|A_\lambda u\| \leq \|u\|, \quad \text{for } u \in K \cap \partial\Omega_{\rho_2}.$$

It follows from Theorem 1.1, that  $A_\lambda$  has a fixed point in  $K \cap (\bar{\Omega}_{\rho_2} \setminus \Omega_{\rho_1})$ , such that  $\rho_1 \leq \|u\| \leq \rho_2$ .  $\square$

From Theorem 3.3, we have

**Corollary 3.4.** *Assume that (H1) and (H2) hold,  $0 < \eta < 1$  and  $0 < \alpha < \frac{1}{\eta}$ . Then we have*

- (1) *If  $f_0 = \infty$ ,  $f_\infty = 0$ , then for each  $\lambda \in (0, \infty)$ , the BVP (1.1) and (1.2) has at least one positive solution.*
- (2) *If  $f_\infty = 0$ ,  $0 < f_0 < \infty$ , then for each  $\lambda \in (\frac{1}{\Lambda_2 f_0}, \infty)$ , the BVP (1.1) and (1.2) has at least one positive solution.*
- (3) *If  $f_0 = \infty$ ,  $0 < f_\infty < \infty$ , then for each  $\lambda \in (0, \frac{1}{\Lambda_1 f_\infty})$ , the BVP (1.1) and (1.2) has at least one positive solution.*

#### 4. EXAMPLES

In this section we present some examples to illustrate our main results.

**Example 4.1.** Consider the boundary value problem

$$(4.1) \quad u''(t) + tu^p = 0, \quad 0 < t < 1,$$

$$(4.2) \quad u'(0) = 0, \quad u(1) = 2 \int_0^{\frac{1}{4}} u(s) ds.$$

Set  $\alpha = 2$ ,  $\eta = 1/4$ ,  $a(t) = t$ ,  $f(u) = u^p$  ( $p \in (0, 1) \cup (1, \infty)$ ). We can show that  $0 < \alpha = 2 < 4 = 1/\eta$ .

Now we consider the existence of positive solutions of the problem (4.1) and (4.2) in two cases.

Case 1:  $p > 1$ . In this case,  $f_0 = 0$ ,  $f_\infty = \infty$ . Then, by Corollary 3.2, the BVP (4.1) and (4.2) has at least one positive solution.

Case 2:  $p \in (0, 1)$ . In this case,  $f_0 = \infty$ ,  $f_\infty = 0$ . Then, by Corollary 3.4, the BVP (4.1) and (4.2) has at least one positive solution.

**Example 4.2.** Consider the boundary value problem

$$(4.3) \quad u''(t) + \frac{au e^{2u}}{b + e^u + e^{2u}} = 0, \quad 0 < t < 1,$$

$$(4.4) \quad u'(0) = 0, \quad u(1) = 2 \int_0^{\frac{1}{3}} u(s) ds.$$

Set  $\alpha = 2$ ,  $\eta = 1/3$ ,  $a(t) \equiv 1$ ,  $f(u) = (au e^{2u})/(b + e^u + e^{2u})$ . we consider the existence of positive solutions of the problem (4.3) and (4.4) into the following two cases.

Case 1: If  $a = 5$ ,  $b = 8$ , then  $f_0 = \frac{1}{2}$ ,  $f_\infty = 5$ . By calculating, it is easy to obtain that  $0 < \alpha = 2 < 3 = 1/\eta$ ,  $\gamma = \frac{2}{3}$ . Again

$$\begin{aligned} \Lambda_1 &= \frac{1}{1 - \alpha\eta} \int_0^1 (1 - s)a(s) ds = \frac{3}{2}, \\ \Lambda_2 &= \frac{\gamma}{2(1 - \alpha\eta)} \int_0^\eta \left(2(1 - \eta) + \alpha(\eta^2 - s^2)\right) a(s) ds = \frac{40}{81}, \\ \Lambda_1 f_0 &= \frac{3}{4}, \quad \Lambda_2 f_\infty = \frac{200}{81}. \end{aligned}$$

By Theorem 3.1, we know that for any  $\lambda \in (\frac{81}{200}, \frac{4}{3})$ , the BVP (4.3) and (4.4) has at least one positive solution  $u \in C[0, 1]$ .

Case 2: If  $a = 5$ ,  $b = -2$ , then  $f_0 = \infty$ ,  $f_\infty = 5$  and  $\Lambda_1 f_\infty = \frac{15}{2}$ .

Therefore, by Corollary 3.4, we know that for any  $\lambda \in (0, \frac{2}{15})$ , the BVP (4.3) and (4.4) has at least one positive solution  $u \in C[0, 1]$ .

**Example 4.3.** Consider the boundary value problem

$$(4.5) \quad u''(t) + \frac{1}{5}u\left(1 - \frac{1}{1 + u^2}\right) = 0, \quad 0 < t < 1,$$

$$(4.6) \quad u'(0) = 0, \quad u(1) = \int_0^{\frac{1}{2}} u(s) ds,$$

where  $\alpha = 1$ ,  $\eta = 1/2$ ,  $a(t) \equiv \frac{1}{5}$ ,  $f(u) = u\left(1 - \frac{1}{1 + u^2}\right)$ . By calculating, we have  $0 < \alpha = 1 < 2 = 1/\eta$ ,  $\gamma = \frac{1}{2}$ ,  $f_0 = 0$ ,  $f_\infty = 1$ . Again

$$\Lambda_2 = \frac{\gamma}{2(1 - \alpha\eta)} \int_0^\eta \left(2(1 - \eta) + \alpha(\eta^2 - s^2)\right) a(s) ds = \frac{7}{120}.$$

Hence, by Corollary 3.2, for any  $\lambda \in (\frac{120}{7}, \infty)$ , the BVP (4.5) and (4.6) has at least one positive solution  $u \in C[0, 1]$ .

**Acknowledgments.** The authors would like to thank the anonymous referee for his/her valuable comments and suggestions on the paper.

#### REFERENCES

- [1] D. R. Anderson, Nonlinear triple-point problems on time scales, *Electron. J. Diff. Eqns.* **47** (2004), 1–12.
- [2] ———, Solutions to second order three-point problems on time scales, *J. Difference Equ. Appl.* **8** (2002), 673–688.
- [3] A. Boucherif, Second-order boundary value problems with integral boundary conditions, *Nonlinear Anal.* **70** (2009), 364–371.
- [4] Z. Chengbo, Positive solutions for semi-positone three-point boundary value problems, *J. Comput. Appl. Math.* **228** (2009), 279–286.
- [5] M. Feng, D. Ji and W. Ge, Positive solutions for a class of boundary-value problem with integral boundary conditions in Banach spaces, *J. Comput. Appl. Math.* **222** (2008), 351–363.
- [6] W. Feng and J. R. L. Webb, Solvability of a three-point nonlinear boundary value problem at resonance, *Nonlinear Anal.* **30** (1997), no. 6, 3227–3238.
- [7] Y. Guo and W. Ge, Positive solutions for three-point boundary value problems with dependence on the first order derivative, *J. Math. Anal. Appl.* **290** (2004), 291–301.
- [8] C. P. Gupta, Solvability of a three-point nonlinear boundary value problem for a second order ordinary differential equations, *J. Math. Anal. Appl.* **168** (1992), 540–551.
- [9] F. Haddouchi and S. Benaicha, Multiple positive solutions for a nonlinear three-point integral boundary-value problem, *Int. J. Open Probl. Comput. Sci. Math.* **8** (2015), no. 1, 29–42.
- [10] X. Han, Positive solutions for a three-point boundary value problem, *Nonlinear Anal.* **66** (2007), 679–688.
- [11] X. He and W. Ge, Triple solutions for second order three-point boundary value problems, *J. Math. Anal. Appl.* **268** (2002), 256–265.
- [12] V. A. Il'in and E. I. Moiseev, Nonlocal boundary-value problem of the first kind for a Sturm-Liouville operator in its differential and finite difference aspects, *Differ. Equ.* **23** (1987), no. 7, 803–810.
- [13] J. Jiang, L. Liu, and Y. Wu, Second-order nonlinear singular Sturm-Liouville problems with integral boundary conditions, *Appl. Math. Comput.* **215** (2009), 1573–1582.
- [14] P. Kang and Z. Wei, Three positive solutions of singular nonlocal boundary value problems for systems of nonlinear second-order ordinary differential equations, *Nonlinear Anal.* **70** (2008), 444–451.
- [15] L. Kong, Second order singular boundary value problems with integral boundary conditions, *Nonlinear Anal.* **72** (2010), 2628–2638.
- [16] M. A. Krasnoselskii, *Positive Solutions of Operator Equations*, P. Noordhoff, Groningen, The Netherlands, 1964.
- [17] J. Li and J. Shen, Multiple positive solutions for a second-order three-point boundary value problem, *Appl. Math. Comput.* **182** (2006), 258–268.
- [18] S. Liang and L. Mu, Multiplicity of positive solutions for singular three-point boundary value problems at resonance, *Nonlinear Anal.* **71** (2009), 2497–2505.
- [19] R. Liang, J. Peng, and J. Shen, Positive solutions to a generalized second order three-point boundary value problem, *Appl. Math. Comput.* **196** (2008), 931–940.
- [20] B. Liu, Positive solutions of a nonlinear three-point boundary value problem, *Appl. Math. Comput.* **132** (2002), 11–28.
- [21] ———, Positive solutions of a nonlinear three-point boundary value problem, *Comput. Math. Appl.* **44** (2002), 201–211.
- [22] H. Luo and Q. Ma, Positive solutions to a generalized second-order three-point boundary-value problem on time scales, *Electron. J. Diff. Eqns.* **17** (2005), 1–14.

- [23] R. Ma, Existence theorems for a second order three-point boundary value problem, *J. Math. Anal. Appl.* **212** (1997), 430–442.
- [24] ———, Multiplicity of positive solutions for second-order three-point boundary value problems, *Comput. Math. Appl.* **40** (2000), 193–204.
- [25] ———, Positive solutions for a nonlinear three-point boundary value problem, *Electron. J. Diff. Eqns.* **34** (1999), 1–8.
- [26] ———, Positive solutions for second-order three-point boundary value problems, *Appl. Math. Lett.* **14** (2001), 1–5.
- [27] R. Ma and H. Wang, Positive solutions of nonlinear three-point boundary value problem, *J. Math. Anal. Appl.* **279** (2003), 216–227.
- [28] Q. Ma, Existence of positive solutions for the symmetry three-point boundary value problem, *Electron. J. Diff. Eqns.* **154** (2007), 1–8.
- [29] S. A. Marano, A remark on a second order three-point boundary value problem, *J. Math. Anal. Appl.* **183** (1994), 581–522.
- [30] H. Pang, M. Feng, and W. Ge, Existence and monotone iteration of positive solutions for a three-point boundary value problem, *Appl. Math. Lett.* **21** (2008), 656–661.
- [31] I. K. Purnaras, On the existence of nonnegative solutions to an integral equation with applications to boundary value problems, *Nonlinear Anal.* **71** (2009), 3914–3933.
- [32] H. R. Sun and W. T. Li, Positive solutions for nonlinear three-point boundary value problems on time scales, *J. Math. Anal. Appl.* **299** (2004), 508–524.
- [33] Y. Sun, L. Liu, J. Zhang, and R. P. Agarwal, Positive solutions of singular three-point boundary value problems for second-order differential equations, *J. Comput. Appl. Math.* **230** (2009), 738–750.
- [34] J. Tariboon and T. Sitthiwirattam, Positive solutions of a nonlinear three-point integral boundary value problem, *Bound. Val. Prob.* ID 519210, doi:10.1155/2010/519210 (2010), 11 pages.
- [35] J. R. L. Webb, Positive solutions of some three-point boundary value problems via fixed point index theory, *Nonlinear Anal.* **47** (2001), no. 7, 4319–4332.
- [36] X. Xu, Multiplicity results for positive solutions of some semi-positone three-point boundary value problems, *J. Math. Anal. Appl.* **291** (2004), 673–689.
- [37] X. Zhang, M. Feng, and W. Ge, Symmetric positive solutions for p-Laplacian fourth-order differential equations with integral boundary conditions, *J. Comput. Appl. Math.* **222** (2008), 561–573.

FACULTY OF PHYSICS, UNIVERSITY OF SCIENCES AND TECHNOLOGY OF ORAN-MB, EL MNAOUAR,  
 BP 1505, 31000 ORAN, ALGERIA  
*E-mail address:* fhaddouchi@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ORAN 1 AHMED BENBELLA, 31000 ORAN, ALGERIA  
*E-mail address:* slimanebenaicha@yahoo.fr